




# Aminoallyl-dUTP - Solid

5-(3-Aminoallyl)-2'-deoxyuridine-5'-triphosphate, Sodium salt

Cat. No.	Amount
NU-803-1	1 mg
NU-803-5	5 mg



Structural formula of Aminoallyl-dUTP - Solid

## For general laboratory use.

Shipping: shipped on gel packs

Storage Conditions: store at -20 °C

Short term exposure (up to 1 week cumulative) to ambient temperature possible.

Shelf Life: 12 months after date of delivery

**Molecular Formula:** C<sub>12</sub>H<sub>20</sub>N<sub>3</sub>O<sub>14</sub>P<sub>3</sub> (free acid)

Molecular Weight: 523.22 g/mol (free acid)

Exact Mass: 523.02 g/mol (free acid)

CAS#: 116840-18-7

**Purity:** ≥ 95 % (HPLC)

Form: solid

Color: white to off-white

Spectroscopic Properties:  $\lambda_{max}$  289 nm,  $\epsilon$  7.1 L mmol<sup>-1</sup> cm<sup>-1</sup> (Tris-HCl pH 7.5)

## **Applications:**

- Incorporation into DNA/cDNA by
  - PCR with Taq polymerase<sup>[1], in-house data</sup>
  - Nick Translation with DNAse I/ DNA Polymerase I<sup>[2]</sup>
  - Primer Extension with Klenow exo<sup>- [3]</sup>
  - 3'-End Labeling with Terminal deoxynucleotidyl Transferase (TdT)<sup>[4]</sup>
  - Reverse Transcription with MMLV Reverse Transcriptase<sup>[2]</sup>

### **Description**:

Aminoallyl-dUTP is recommended for two-step labeling of DNA/cDNA e.g. by PCR, Nick Translation, Primer Extension, 3'-End Labeling and Reverse Transcription. It is enzymatically incorporated into DNA/cDNA as substitute for its natural counterpart dTTP. The resulting Amine-functionalized DNA/cDNA can subsequently be labeled via the classic Amine/NHS Ester reaction that offers the choice

- to introduce a Biotin group (via NHS Ester of Biotin) for subsequent purification tasks
- to introduce fluorescent group (via NHS Ester of fluorescent dyes) for subsequent microscopic imaging

#### Selected References:

[1] Dirsch *et al.* (2007) Probe production for *in situ* hybridization by PCR and subsequent covalent labeling with fluorescent dyes. *Appl. Immunohistochem. Mol. Morphol.* **3**:332.

[2] Cox *et al.* (2004) Fluorescent DNA hybridization probe preparation using amine modification and reactive dye coupling. *BioTechniques* **36**:114.

[3] Cherkasov *et al.* (2010) New Nucleotide Analogues with Enhanced Signal Properties. *Bioconjugate Chem.* **21 (1)**:122.

[4] Unciti-Broceta *et al.* (2003) The use of solid supports to generate nucleic acid carriers.*Accounts of Chemical Research* **45**:1140.

